

Actinium
Pharmaceuticals, Inc.

Administration of ^{131}I -apamistamab

ICD-10-PCS Coordination & Maintenance Committee Meeting
September 2023

Disclaimer and Safe Harbor

The information presented herein contains express and implied forward-looking statements regarding the current intentions, expectations, estimates, opinions and beliefs of Actinium Pharmaceuticals, Inc. (“Actinium”) that are not historical facts. These forward-looking statements include statements regarding Actinium’s expectations for its product candidates (including their therapeutic and commercial potential, anticipated future development activities, anticipated timing of development activities, including initiation of clinical trials and presentations of clinical data and the indications Actinium and its collaborators plan to pursue), future results of operations and financial position, business strategy, strategic collaborations, any royalty or milestone payments and Actinium’s ability to obtain and maintain intellectual property protection for its product candidates. Such forward-looking statements may be identified by words such as “believes”, “may”, “will”, “expects”, “endeavors”, “anticipates”, “intends”, “plans”, “estimates”, “projects”, “should”, “objective” and variations of such words and similar words. These statements are based on management’s current expectations and are subject to risks and uncertainties that may cause actual results to differ materially from the anticipated or estimated future results, including the risks and uncertainties associated with preliminary study results varying from final results, estimates of potential markets for drugs under development, clinical trials, actions by the FDA and other governmental agencies, regulatory clearances, responses to regulatory matters, the market demand for and acceptance of Actinium’s products and services, performance of clinical research organizations and other risks detailed from time to time in Actinium’s filings with the Securities and Exchange Commission (the “SEC”), including without limitation its most recent annual report on Form 10-K, subsequent quarterly reports on Forms 10-Q and Forms 8-K, each as amended and supplemented from time to time.

Any forward-looking statements that Actinium makes in this presentation speak only as of the date of this presentation. Except as required by law, Actinium assumes no obligation to update its forward-looking statements whether as a result of new information, future events or otherwise, after the date hereof. Nothing contained in this presentation is, or should be construed as, a recommendation, promise or representation by Actinium or any director, employee, agent, or adviser of Actinium. This presentation does not purport to be all-inclusive or to contain all of the information you may desire. The content of this presentation is subject to copyright, which will be asserted by Actinium, and no part of this presentation may be reproduced, stored in a retrieval system, or transmitted in any form or by any means without prior permission in writing from Actinium.

Acute Myeloid Leukemia (AML) Fast Facts

AML is an aggressive disease that can progress rapidly despite treatments

≈21,000
AML patients
annually¹


68 years
Median age at
diagnosis¹

10 drugs
approved for
AML patients
since 2017

Older patients have
limited treatment
options and poor
outcomes



Develop R/R
disease²



of AML patients
access HCT³

HCT is the only
potential curative
treatment
for R/R AML,
but only younger or fit
patients can access it


Challenges to Achieving Cures in AML

Patients must be able to overcome several challenges related to curative allogeneic hematopoietic cell transplant (alloHCT)

- | | | |
|----------------|--|----------|
| ➡ Challenge #1 | Need to attain a complete remission (CR) | Access |
| ➡ Challenge #2 | Tolerate and survive effective alloHCT conditioning | |
| ➡ Challenge #3 | <ul style="list-style-type: none">• Achieve alloHCT engraftment• Achieve post-HCT CR | Outcomes |
| ➡ Challenge #4 | <ul style="list-style-type: none">• Surmount alloHCT related complications<ul style="list-style-type: none">– Graft failure– Side effects: sepsis, GVHD | |

Overcoming these challenges can result in long-term survival and curative outcomes.

¹³¹I-apamistamab: A Next Generation Approach to Improve alloHCT Access, Outcomes




- ¹³¹I-apamistamab targets CD45, which is highly expressed on AML, immune and stem cells
- Low CD45 expression outside the hematopoietic system spares organs and increases tolerability of ¹³¹I-apamistamab
- Enables high amounts of radiation to be delivered to radiation sensitive AML and immune cells
- Induction and conditioning in one approach by simultaneously eliminating AML immune and bone marrow stem cells
- Allows patients with active disease to go directly to alloHCT rapidly via a single infusion


¹³¹I-apamistamab targeted radiotherapy can produce myeloablative outcomes with the safety and tolerability of reduced intensity approaches

Personalized Single Dose Combined Induction/Conditioning

- This shows a single inpatient stay, where the patient receives the therapeutic dose (dosimetric dosing will typically occur on an outpatient basis). The patient then proceeds to a reduced intensity conditioning and to allogeneic hematopoietic cell transplant (alloHCT).



- This shows two inpatient stays; one where the patient receives the therapeutic dose (and is isolated until an acceptable level of radiation is achieved), and then receives fludarabine for days -4 to -2, and is admitted as an inpatient on day -0 for TBI and alloHCT.




Dosage and Documentation

- The administration of the therapeutic dose for ^{131}I -apamistamab will be documented in the medical record in the same manner as infusion of other targeted therapy and in the treating bone marrow transplanter (hematologist/oncologist) physician's note
- The full name for ^{131}I -apamistamab is ^{131}I iodine apamistamab (also known as lomab-B)
 - It is also described as Radiolabeled ^{131}I iodine Anti-CD45 (BC8) Antibody in scientific literature


SIERRA Results: ^{131}I -apamistamab Overcomes Key HCT Challenges

Primary endpoint met with high statistical significance: High rates of post-HCT remissions resulted in significantly higher durable complete remissions with ^{131}I -apamistamab




Challenge 1 & 2 Addressed: Unprecedented Access to HCT in Half the Time

Unprecedented Access to HCT




- **Targeted nature of ¹³¹I-apamistamab** results in effective myeloablation with improved tolerability
- **Less than 20%** of patients on the control arm had access to transplant vs. 100% on ¹³¹I-apamistamab

Access to HCT in Less Than Half the Time



- **With ¹³¹I-apamistamab there is no need to achieve a CR;** patients with active disease can go to directly to BMT in days and engraft successfully
- **1 – 2+ cycles of chemotherapy** required to attain CR if CR can even be produced

Challenge 3 Addressed: Unprecedented HCT Access & Engraftment and High Post-HCT CR



Challenge 4 Addressed: Excellent Safety of Targeted Radiotherapy

¹³¹I-apamistamab related adverse events are meaningfully lower, implying less complexity (and cost) pre- and post-transplant.

Adverse Event* (%)	¹³¹ I-apamistamab Arm N=66	Control Arm N=14
Sepsis ¹	6.1%	28.6%
Febrile Neutropenia	43.9%	50.0%
Mucositis ²	15.2%	21.4%
Acute GVHD (Gr II-IV) ³	26.1%	35.7%

* Relevant adverse events in transplanted ¹³¹I-apamistamab patients. 1) "Sepsis" includes Preferred Terms of Sepsis, Septic Shock, Neutropenic Sepsis & Septic Embolus; 2) "Mucositis" includes Preferred Terms of Stomatitis & Mucosal Inflammation; 3) All ¹³¹I-apamistamab pts received Cyclosporin and Mycophenolate Mofetil for GVHD prophylaxis. Late Breaking Abstract, TCT 2023, Efficacy and Safety Results of the Sierra Trial: A Multicenter, Pivotal Phase 3 Study of Iomab-B Prior to Allogeneic HCT Versus Conventional Care in Older Patients with Active, R/R AML


Challenge 4 Addressed: Excellent Safety of Targeted Radiotherapy (cont.)

This table shows supplemental information on the \geq grade 3 common drug-related Adverse Events (AEs) in $\geq 5\%$ in either group.


	Crossed Over to lomab-B (N=44), N (%)	lomab-B Arm (N=72), N (%)	Total (N=116), N (%)
Blood & Lymphatic System Disorders	14 (31.8)	21 (29.2)	35 (30.2)
Investigations	8 (18.2)	18 (25.0)	26 (22.4)
Infections & Infestations	8 (18.2)	14 (19.4)	22 (19.0)
Metabolism & Nutrition Disorders	8 (18.2)	6 (8.3)	14 (12.1)
Gastrointestinal Disorders	6 (13.6)	7 (9.7)	13 (11.2)
Respiratory, Thoracic & Mediastinal Disorders	3 (6.8)	9 (12.5)	12 (10.3)
General Disorders & Administration Site Conditions	4 (9.1)	7 (9.7)	11 (9.5)

^{131}I -apamistamab Demonstrates Clear Survival Benefit

^{131}I -apamistamab doubled 1-year survival rates and median overall survival



¹³¹I-apamistamab Reduced the Probability of an Event by 78%



EFS HR of 0.22 (P<0.0001) compared to control arm clearly supports the use of ¹³¹I-apamistamab in this high-risk population

^{131}I -apamistamab Represents a New Paradigm

Clear separation in access and outcomes favor ^{131}I -apamistamab utilization

